

A new genus of prosopid crabs (Decapoda: Brachyura: Dromiacea) from the Upper Jurassic of southern Germany

Un nuevo género de cangrejos prosópidos (Decapoda: Brachyura: Dromiacea) del Jurásico Superior del sur de Alemania

Günter Schweigert^{1,*}

¹ Staatliches Museum für Naturkunde, Rosenstein 1, 70191 Stuttgart, Germany.

* Corresponding author: (G. Schweigert)
guenter.schweigert@smns-bw.de

ABSTRACT

A new monotypic genus and species of brachyurans, *Petersbuchia thaukei* n. gen. n. sp., is described from southern Germany. The holotype, an incompletely preserved carapace, was collected from a thick-bedded spongolithic limestone of the Upper Jurassic Treuchtlingen Formation (early late Kimmeridgian, Acanthicum Zone). *Petersbuchia* adds to the diversification hotspot of early true crabs within the sponge-microbial magnafacies along the northern shelf of the Tethys Ocean in general and to the brachyuran fauna of the Treuchtlingen Formation in special.

Keywords: Homolodromioidea, diversification, Late Jurassic, palaeoecology, Treuchtlingen Formation, Bavaria, Germany.

RESUMEN

Un nuevo género y especie monotípico de brachyuros, *Petersbuchia thaukei* n. gen. n. sp., es descrito del sur de Alemania. El holotipo, un caparazón incompleto, fue colectado de una gruesa capa espongiolítica de la Formación Treuchtlingen del Jurásico Superior (Kimmeridgiano temprano tardío, Zona Acanthicum). *Petersbuchia* se añade el punto de diversificación de los cangrejos verdaderos en magnafacies espongío-microbianas a lo largo de la Plataforma norte del Océano del Tethys en general, y de la fauna de braquiuros de la Formación Treuchtlingen, en particular.

Palabras clave: Homolodromioidea, diversificación, Jurásico Tardío, palaeoecología, Formación Treuchtlingen, Bavaria, Alemania.

How to cite this article:

Schweigert, G., 2021, A new genus of prosopid crabs (Decapoda: Brachyura: Dromiacea) from the Upper Jurassic of southern Germany: Boletín de la Sociedad Geológica Mexicana, 73 (3), A030121. <http://dx.doi.org/10.18268/BSGM2021v73n3a030121>

Manuscript received: November 30, 2020
Corrected manuscript received: December 26, 2020
Manuscript accepted: January 4, 2021

Peer Reviewing under the responsibility of
Universidad Nacional Autónoma de México.

This is an open access article under the CC BY-NC-SA license (<https://creativecommons.org/licenses/by-nc-sa/4.0/>)

1. Introduction

In modern marine benthic environments, brachyurans are well represented by approximately 7,250 species (Davie *et al.*, 2015). The most primitive ones, sometimes controversially discussed, have been recorded from clayey marine strata of Early Jurassic (Pliensbachian) age (Förster, 1986; Schweitzer and Feldmann, 2010d; Haug and Haug, 2014; Krzemińska *et al.*, 2020; Scholz, 2020). Brachyuran records from the Middle Jurassic are still scarce (Krobicki and Zatoń, 2008; Fraaije *et al.*, 2013). In the Late Jurassic, siliceous sponge-microbial magnafacies as well as coeval coral reefs became widespread along the northern margin of the Tethys Ocean (e.g., Leinfelder *et al.*, 1994, 1996). This probably triggered the first great diversification and biogeographical expansion of crabs (Wehner, 1988; Müller *et al.*, 2000; Klompmaker *et al.*, 2013). The Upper Jurassic of southern Germany, which was part of this northern Tethyan reef belt, is not only a classical area for early brachyuran studies (von Meyer, 1842, 1860) but was probably also a diversification hotspot of this group, although geographically isolated brachyuran records even come from the western Pacific area (Karasawa and Kato, 2007; Kato *et al.*, 2010; Karasawa and Hirota, 2019). Early records from the southern Tethys (Förster, 1985) are of dubious origination (see Krobicki and Zatoń, 2008).

In the course of an extensive revision and update of the 'Treatise' volume dedicated to decapod crustaceans (Glaessner, 1969), numerous taxonomic revisions of the original type material from old collections became necessary, and additional taxa based on newly discovered or differently interpreted Jurassic crab material were occasionally added (Feldmann *et al.*, 2006; Schweitzer *et al.*, 2007, 2018; Schweitzer and Feldmann, 2008a, 2008b, 2009a, 2009b, 2009c, 2009d; 2010a, 2010b, 2010c; Frantescu, 2011; Schweigert and Koppka, 2011; Starzyk *et al.*, 2012; Hyžný *et al.*, 2015; Starzyk, 2015a, 2015b, 2020; Schweigert and Robins, 2016; Schweigert, 2019; Klompmaker *et al.*, 2020).

The aim of this study is to report a new genus and species of the Prosopidae. The abbreviation SMNS is used for Staatliches Museum für Naturkunde Stuttgart, Germany.

2. Geological setting

The herein described specimen was collected from thick-bedded limestones of the Treuchtlingen Formation (Niebuhr and Pürner, 2014) quarried in the Petersbuch-Erkertshofen district, which is part of the township of Markt Titting in Bavaria (Germany) (Figure 1). Due to its usage as a building stone these limestones are well-known and traded as "Treuchtlingen Marble". The limestones of the Treuchtlingen Formation yield abundant ammonites, belemnite rostra, siliceous and calcareous sponges, brachiopods, and echinoids. In contrast, hermatypic corals are totally absent from this lithology. In the finer rock matrix macroscopically visible foraminifera with a whitish wall ("Tubiphytes") are extremely common and characteristic for this lithology. The ammonite fauna allows a precise dating of these rocks within the late Kimmeridgian (Zeiss, 1964; Bantz, 1970; Schlampp, 2004, 2009). The environmental conditions during which the Treuchtlingen Formation was deposited have been discussed controversially in the past. A very shallow depositional model was suggested by Kott (1989); however, subsequently many arguments, especially the total absence of a typical shallow-water fauna (hermatypic corals, thick-shelled bivalves, large gastropods), have been put forward in favour of greater water depths within a range of ca. 50 to 150 metres, certainly below the storm-weather wave basis (e.g., Leinfelder *et al.*, 1994, 1996; Schmid, 1996; Schmid *et al.*, 2005). The occurring decapod crustacean fauna consists of small-sized crabs (see below), is relatively low-diverse and does not comprise any species otherwise restricted to Upper Jurassic coral reefs (cf. Schweitzer *et al.*, 2018).

Although numerous amateur collectors are hunting for fossils in the Franconian Treuchtlingen Formation these tiny fossils are not in their focus

and they are mostly ignored and overlooked. Here is a list of brachyurans from the Treuchtlingen Formation (compiled after Wehner, 1988, Schweigert and Kuschel, 2018, and own observations; taxonomy modified after Schweitzer *et al.*, 2018):

Eodromites grandis (von Meyer, 1857)
Laeviprosopon laeve (von Meyer, 1857)

Nodoprosopon ornatum (von Meyer, 1842)
Petersbuchia thaukei n. gen., n. sp. (herein)
Pilidromia thiedeae (Schweigert and Koppka, 2011)
Planoprosopon heydeni (von Meyer, 1857)
Tanidromites insignis (von Meyer, 1857)
Tanidromites scheffnerae Schweigert and Koppka, 2011.
Verrucarcinus torosus (von Meyer, 1857)




Figure 1 Locality of the holotype. Arrow indicates location of the Petersbuch quarry at Markt Titting-Petersbuch, Bavaria, southern Germany. Map based on OpenStreetMap (www.openstreetmap.org).

3. Systematic palaeontology

The classification follows Schweitzer *et al.* (2010).

Order Decapoda Latreille, 1802
 Infraorder Brachyura Linnaeus, 1758
 Section Dromiacea De Haan, 1833
 Superfamily Homolodromioidea Alcock, 1900
 Family Prosopidae von Meyer, 1860
 Genus *Petersbuchia* nov.

Type species: *Petersbuchia thaukei* n. gen., n. sp., monotypic. Gender is feminine.

Diagnosis: See diagnosis of the type species (monotypic).

Petersbuchia thaukei n. gen., n. sp.

Figures 2A to 2D, 3

Etymology: Named after Thomas Haucke (Freystadt-Mörsdorf), who collected and donated the sole specimen.

Holotype: Specimen SMNS 70622 (Figure 2).

Type locality: Quarry area of Petersbuch/Erkertshofen, Markt Titting, Franconian Alb, Germany (Figure 1).

Type horizon: Treuchtlingen Formation, Petersbuch Member (Upper Jurassic, late Kimmeridgian, Acanthicum Zone; Niebuhr and Pürner, 2014). Exact bed not traceable, because the rock with the fossil was found ex-situ on top of a dump close to a quarry.

Studied material: Holotype only.

Occurrence: At present only known from the type locality.

Diagnosis: Carapace subquadrate, with long protrusive outer orbital spine; Surface coarsely pustulose, anterior part with a few prominent tubercles. Rostrum trifid, with spatulate margins. Hepatic and gastric areas not separated by grooves.

Description: The small-sized holotype SMNS 70622 (Figure 2) is represented by the three-dimensionally preserved inner cast of the carapace. The left side of the carapace is partly lost prior to burial due to breakage. Surface of

carapace densely pustulose in all areas. Outline of carapace subquadrate, with broad trifid rostrum. Total length of carapace excluding rostrum 3.7 mm, including rostrum 4.6 mm; reconstructed width by mirroring in symmetry plane 3.6 mm, maximum height 2.1 mm. Trifid rostrum with spatulate upturned margins and downturned tip in the centre. Rounded orbital margin connects base of rostrum and protrusive outer orbital spine. Frontal aspects of orbital area hidden. Hepatic, protogastric and epigastric areas not separated from each other. Mesogastric area well separated by lateral grooves, with two coarser tubercles in its symmetry plane. Posterior gastric muscle scars with subparallel, spring-like ornamentation. In addition to the general pustulose ornamentation, four strong and prominent tubercles are developed in the anterior part of the carapace, one close to the mesogastric groove, another one in the outer third of the hepatic region, close to the cervical groove, the third one near the centre of the joint hepatic-protogastric region and the last one in the epigastric region, at the base of the rostrum. Subhepatic region not separated from hepatic region but posteriorwards forming a lobe. Cervical groove and branchiocardioc groove well developed. Epi- and mesobranchial regions fused. Meta- and urogastric regions fused as well. Cardiac region triangular, but hardly separated from adjacent branchial region. Outer lateral margin of branchial region parallel to longitudinal axis of carapace. Posterior margin of carapace forming a smooth rim, with concavities placed laterally from the intestinal area. Sternal region not preserved. Close to the specimen, there is an incompletely preserved chela possibly belonging to the same individual (Figure 3). The propodus is twice as long as broad and widens distally. Surface of propodus with similar pustulose ornamentation as in the carapace. Fixed finger thick, smooth, distally broken, but slightly curved imprint points to a sickle-like outline. Free finger lost.

Remarks: It is impossible to state whether the sole small-sized specimen of *Petersbuchia thaukei* n. gen., n. sp. was an adult or a juvenile.



Figure 2 *Petersbuchia thaukei* n. gen. n. sp., holotype, SMNS 70622, Upper Jurassic, Treuchtlingen Formation, Petersbuch Member; late Kimmeridgian, Acanthicum Zone, Markt Titting-Petersbuch, Bavaria, southern Germany. Dorsal (A, C), lateral (B) and frontal (D) views. A with artificially enhanced contrast. Scale bar = 2 mm.



Figure 3 *Petersbuchia thauckei* n. gen. n. sp., holotype, SMNS 70622, with a fragmentary chela possibly belonging to the same individual (inset, close-up view). Scale bar = 2 mm.

4. Discussion

At present, *Petersbuchia thauckei* n. gen. n. sp. from the late Kimmeridgian of Franconia represents the only species assigned to this new genus. In first view, the number and arrangement of prominent tubercles in the hepatic region of *Petersbuchia thauckei* n. gen., n. sp. is strikingly reminiscent to *Laeviprosopon sublaeve* (von Meyer, 1857), another very rare taxon reported from Upper Jurassic beds of similar age and lithology.

Some species of *Laeviprosopon* including the type species exhibit a similar trifid rostrum as in *Petersbuchia* n. gen. However, *Laeviprosopon sublaeve* lacks a pustulose ornamentation of the carapace as well as an outer orbital spine. It must be stressed that ornamentation is usually less developed in inner

casts than on the cuticle. Therefore, if the cuticle of *Petersbuchia* n. gen. was preserved it would have been even stronger ornamented. Moreover, in *Laeviprosopon sublaeve* the anterior part of the carapace exhibits a trapezoidal outline.

The posterior margin of the carapace showing laterally placed concavities is shared with a few other genera safely assigned to Prosopidae (Prosopon von Meyer, 1835 s. str.; *Europrosopon* Klompmaker, Starzyk, Fraaije and Schweigert, 2020; *Protuberosa* Schweitzer and Feldmann, 2009a), but this character does not occur within species of *Laeviprosopon* (see Klompmaker *et al.*, 2020). Hence, *Petersbuchia* n. gen. is well distinct from *Laeviprosopon* and shows only some superficially convergent characters with this co-occurring genus.

Acknowledgements

I thank Thomas Haucke (Freystadt-Mörsdorf, Germany), who kindly donated the studied specimen and Victor Schlampp (Rednitzhembach, Germany), who provided insightful biostratigraphical information of the Treuchtlingen Formation. Finally, I thank Alessandro Garassino (Deptment of Biology and Earth Sciences, Loma Linda University, USA) and Francisco Vega (Instituto de Geología, Universidad Nacional Autónoma de México, Mexico) for their careful reviews.

References

- Alcock, A., 1900, Materials for a carcinological fauna of India, No. 5. The Brachyura Primigenia or Dromiacea: Journal of the Asiatic Society of Bengal, 68, 123-169. <https://doi.org/10.5962/bhl.title.15343>
- Bantz, H.U., 1970, Der Fossilinhalt des Treuchtlinger Marmors (Mittleres Unter-Kimmeridge der Südlichen Frankenalb): Erlanger Geologische Abhandlungen, 82, 1-86.
- Davie, P.J.F., Guinot, D., Ng, P.K.L., 2015, Systematics and classification of Brachyura, in Castro, P., Davie, P.J.F., Guinot, D., Schram, F. R., von Vaupel Klein, J.C. (eds.), Treatise on zoology – anatomy, taxonomy, biology. The Crustacea, 9C-II: Leiden, Brill, 1049-1130. https://doi.org/10.1163/9789004190832_021
- De Haan, W., 1833, Crustacea, in von Siebold, P. F. (ed.), Fauna Japonica sive Descriptio Animalium, quae in Itinere per Japoniam, Jussu et Auspiciis Superiorum, qui summum in India Batava Imperium Tenent, Suscepto, Annis 1823-1830 Collegit, Notis, Observationibus et Adumbrationibus Illustravit. J. Müller et Co., Lugduni Batavorum, 243 p. <https://doi.org/10.5962/bhl.title.124951>
- Feldmann, R.M., Lazăr, I., Schweitzer, C.E., 2006, New crabs (Decapoda: Brachyura: Prosopidae) from Jurassic (Oxfordian) sponge bioherms of Dobrogea, Romania: Bulletin of the Mizunami Fossil Museum, 33, 1-20.
- Förster, R., 1985, Frühe Anomuren und Brachyuren (Decapoda, Crustacea) aus dem mittleren Dogger: Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie, 25, 45-60.
- Förster, R., 1986, Der erste Nachweis eines brachyuren Krebses aus dem Lias (oberes Pliensbach) Mitteleuropas: Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie, 26, 25-31.
- Fraaije, R.H.B., van Bakel, B.W.M., Guinot, D., Jagt, J.W.M., 2013, A new Middle Jurassic (Bajocian) homolodromiid crab from northwest France; the earliest record of the Tanidromitidae: Boletín de la Sociedad Geológica Mexicana, 65, 249-254. <https://doi.org/10.18268/bsgm2013v65n2a6>
- Franțescu, O. D., 2011, Brachyuran decapods (including five new species and one new genus) from Jurassic (Oxfordian-Kimmeridgian) coral reef limestones from Dobrogea, Romania: Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, 259, 271-297. <https://doi.org/10.1127/0077-7749/2010/0110>
- Glaessner, M., 1969, Decapoda, in Moore, R.C. (ed.), Treatise on Invertebrate Paleontology, Pt. R4(2). Boulder and Lawrence, Geological Society of America and University of Kansas Press, R400-R533, R626-628.
- Haug, J.T., Haug, C., 2014, Eoprosopon klugi (Brachyura) the oldest unequivocal and most “primitive” crab reconsidered: Palaeodiversity, 7, 149-158.
- Hyžný, M., Starzyk, N., Robins, C., Kočová Veselská, M., 2015, Taxonomy and palaeoecology of a decapod crustacean assemblage from the Oxfordian of Stránská skála (Southern Moravia, Czech Republic):

- Bulletin of Geosciences, 90, 633-650. <https://doi.org/10.3140/bull.geosci.1559>
- Karasawa, H., Hirota, T., 2019, Two new species of Planoprosopon (Decapoda: Brachyura: Longodromitidae) from the Torinosu Group (Upper Jurassic–Lower Cretaceous), Shikoku, Japan: Bulletin of the Mizunami Fossil Museum, 45, 27-32.
- Karasawa, H., Kato, H., 2007, New prosopid crabs (Crustacea, Decapoda, Brachyura) from the Upper Jurassic Torinosu Group, Shikoku, Japan: Memorie della Società italiana di Scienze naturali e del Museo civico di Storia naturale di Milano, 35, 62-65.
- Kato, H., Takahashi, T., Taira, M., 2010, Late Jurassic decapod crustaceans from northeast Japan: Palaeontology, 53(4), 761-770. <https://doi.org/10.1111/j.1475-4983.2010.00960.x>
- Klompmaker, A.A., Schweitzer, C.E., Feldmann, R.M., Kowalewski, M., 2013, The influence of reefs on the rise of Mesozoic marine crustaceans: Geology, 41, 1179-1182. <https://doi.org/10.1130/G34768.1>
- Klompmaker, A.A., Starzyk, N., Fraaije, R.H.B., Schweigert, G., 2020, Systematics and convergent evolution of reef-associated Jurassic and Cretaceous crabs (Decapoda, Brachyura): Palaeontologica Electronica, 23(2), a32. <https://doi.org/10.26879/1045>
- Kott, R., 1989, Fazies und Geochemie des Treuchtlinger Marmors: Unter- und Mittel-Kimmeridge, Südliche Frankenalb: Berliner Geowissenschaftliche Abhandlungen, Reihe a Geology Palaeontology, 111, 1-115.
- Krobicki, M., Zatoń, M., 2008, Middle and Late Jurassic roots of brachyuran crabs: palaeoenvironmental distribution during their early evolution: Palaeogeography, Palaeoclimatology, Palaeoecology, 263, 30-43. <https://doi.org/10.1016/j.palaeo.2008.01.025>
- Krzemińska, E., Starzyk, N., Schweigert, G., Whicher, J., Chandler, R.B., Fraaije, R.H.B., 2020, New records of Eogastrodorus (Decapoda, Anomura, Gastrodoridae) from the Middle Jurassic (Bajocian) of England and France: Neues Jahrbuch für Geologie und Paläontologie Abhandlungen, 296(1-2), 147-156. <https://doi.org/10.1127/njgpa/2020/0876>
- Latreille, P.A., 1802, Histoire naturelle, générale et particulière, des Crustacés et des Insectes, v. 3. F. Dufart, Paris.
- Leinfelder, R.R., Krautter, M., Laternser, R., Nose, M., Schmid, D.U., Schweigert, G., Werner, W., Keupp, H., Brugger, H., Herrmann, R., Rehfeld-Kiefer, U., Schroeder, J.H., Reinhold, C., Koch, R., Zeiss, A., Schweizer, V., Christmann, H., Menges, G., Luterbacher, H., 1994, The origin of Jurassic reefs. Current research developments and results: Facies, 31, 1-56. <https://doi.org/10.1007/BF02536932>
- Leinfelder, R. R., Werner, W., Nose, M., Schmid, D. U., Krautter, M., Laternser, R., Takać, M., Hartmann, D., 1996, Paleoecology, growth parameters and dynamics of coral, sponge and microbolite reefs from the Late Jurassic. In Reitner, J., Neuweiler, F., Gunkel, F. (eds.), Global and regional controls on biogenic sedimentation. I. Reef evolution. Research Reports: Göttinger Arbeiten zur Geologie und Paläontologie, Sonderband, 2, 227-248.
- Linnaeus, C., 1758, Systema Naturae per Regna tria Naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis (ed. 10), vol 1. Laurentius Salvius, Holmiae. <https://doi.org/10.5962/bhl.title.156772>
- Müller, P., Krobicki, M., Wehner, G., 2000, Jurassic and Cretaceous primitive crabs of the family Prosopidae (Decapoda: Brachyura)-their taxonomy, ecology and biogeography: Annales Societatis Geologorum Poloniae, 70, 49-79.
- Niebuhr, B., Pürner, T., 2014, Plattenkalk und Frankendolomit - Lithostratigraphie der Weißjura-Gruppe der Frankenalb (außerälpiner Oberjura, Bayern):

- Schriftenreihe der Deutschen Gesellschaft für Geowissenschaften, 83, 5-71. <https://doi.org/10.1127/sdgg/83/2014/5>
- Scholz, G., 2020, *Eocarcinus praecursor* Withers, 1932 (Malacostraca, Decapoda, Meiura) is a stem group brachyuran: Arthropod Structure & Development, 59, 100991. <https://doi.org/10.1016/j.asd.2020.100991>
- Schlamp, V., 2004, Die Malm-Ammoniten der Fränkischen Alb gamma 3 und delta: Fossilien, 21(4), 208-219.
- Schlamp, V., 2009, Ammoniten aus dem Treuchtlinger Marmor: Der Steinkern, 3, 40-48.
- Schmid, D. U., 1996, Marine Mikrobolithe und Mikroinkrustierer aus dem Oberjura: Profil, 9, 101-251.
- Schmid, D. U., Leinfelder, R. R., Schweigert, G., 2005, Stratigraphy and palaeoenvironments of the Upper Jurassic of Southern Germany – a review: *Zitteliana*, 26B, 31-41.
- Schweigert, G., 2019, An additional representative of Lecythocaridae (Brachyura: Dromiacea) from the Upper Jurassic of southern Germany: *Journal of Crustacean Biology*, 39(1), 36-39. <https://doi.org/10.1093/jcobiol/ruy092>
- Schweigert, G., Koppka, J., 2011, Decapods (Crustacea: Brachyura) from the Jurassic of Germany and Lithuania, with descriptions of new species of Planoprosopon and Tanidromites: *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 260, 221-235. <https://doi.org/10.1127/0077-7749/2011/0136>
- Schweigert, G., Kuschel, H., 2018, Schwäbische und Fränkische Alb – ein Paradies für Jura-Krabben: Fossilien, 35(4), 14-23.
- Schweigert, G., Robins, C., 2016, Earliest representatives of Lecythocaridae (Crustacea: Brachyura) from the Upper Jurassic of southern Germany. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 280, 325-330. <https://doi.org/10.1127/njgpa/2016/0583>
- Schweitzer, C.E., Feldmann, R.M., 2008a, A new classification for some Jurassic Brachyura (Crustacea: Decapoda: Brachyura: Homolodromioidea): Families Goniodromitidae Beurlen, 1932 and Tanidromitidae new family: *Senckenbergiana lethaea*, 87, 119-156. <https://doi.org/10.1007/BF03043911>
- Schweitzer, C.E., Feldmann, R.M., 2008b, Revision of the genus *Laeviprosopon* Glaessner, 1933, (Decapoda: Brachyura: Homolodromioidea: Prosopidae) including two new species: *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 250, 273-285. <https://doi.org/10.1127/0077-7749/2008/0250-0273>
- Schweitzer, C.E., Feldmann, R.M., 2009a, Revision of the Prosopinae sensu Glaessner, 1969 (Crustacea: Decapoda: Brachyura) including four new families, four new genera, and five new species: *Annalen des Naturhistorischen Museums in Wien*, 110A, 55-121.
- Schweitzer, C.E., Feldmann, R.M., 2009b, New species of Longodromitidae Schweitzer and Feldmann, 2009, from the Ernstbrunn Formation, Late Jurassic (Tithonian), Austria: *Annalen des Naturhistorischen Museums in Wien*, 111A, 207-224.
- Schweitzer, C.E., Feldmann, R.M., 2009c, Revision of Gabriella Collins *et al.*, 2006 (Decapoda: Brachyura: Homolodromioidea: Tanidromitidae) with new Jurassic species: *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 252, 1-16. <https://doi.org/10.1127/0077-7749/2009/0252-0001>
- Schweitzer, C.E., Feldmann, R.M., 2009d, Revision of the genus *Cyclothyreus* Remeš, 1895 (Decapoda: Brachyura: Dromioidea): *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 253, 357-372. <https://doi.org/10.1127/0077-7749/2009/0253-0357>
- Schweitzer, C.E., Feldmann, R.M., 2010a, A new family of Mesozoic Brachyura (Glaessneropsoidae) and reevaluation of *Distefania* Checchia-Rispoli, 1917

- (Homolodromoidea: Goniodromitidae): Neues Jahrbuch für Geologie und Paläontologie Abhandlungen, 256, 363-380. <https://doi.org/10.1127/0077-7749/2010/0059>
- Schweitzer, C.E., Feldmann, R.M., 2010b, Revision of Cycloprosopon and additional notes on Eodromites (Brachyura: Homolodromoidea: Goniodromitidae): Annalen des Naturhistorischen Museums in Wien, Serie A, 112, 169-194.
- Schweitzer, C.E., Feldmann, R.M., 2010c, The genus Coelopus Étallon, 1861 (Brachyura: Glaessneropsoidea: Longodromitidae) with new species: Neues Jahrbuch für Geologie und Paläontologie Abhandlungen, 258, 51-60. <https://doi.org/10.1127/0077-7749/2010/0084>
- Schweitzer, C.E., Feldmann, R.M., 2010d, The oldest Brachyura (Decapoda: Homolodromoidea: Glaessneropsoidea) known to date (Jurassic): Journal of Crustacean Biology, 30, 251-256. <https://doi.org/10.1651/09-3231.1>
- Schweitzer, C.E., Feldmann, R.M., Garassino, A., Karasawa, H., Schweigert, G., 2010, Systematic list of fossil crustacean species: Crustaceana Monographs, 10, 222 p. <https://doi.org/10.1163/ej.9789004178915.i-222>
- Schweitzer, C.E., Feldmann, R.M., Lazăr, I., 2007, Decapods from Jurassic (Oxfordian) sponge megafacies of Dobrogea, Romania and reconsideration of Nodoprosopon Beurlen, 1928: Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, 244, 99-113. <https://doi.org/10.1127/0077-7749/2007/0244-0099>
- Schweitzer, C.E., Feldmann, R.M., Lazăr, I., Schweigert, G., Franțescu, O., 2018, Decapoda (Anomura; Brachyura) from the Late Jurassic of the Carpathians, Romania: Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, 288, 307-341. <https://doi.org/10.1127/njgpa/2018/0744>
- Starzyk, N., 2015a, Reappraisal of the primitive crab Eodromites, with description of three new species from European localities (Decapoda: Brachyura: Goniodromitidae): Palaeontologica Electronica, 18.3.50A, 1-19. <https://doi.org/10.26879/513>
- Starzyk, N., 2015b, New species of the genus Tanidromites (Decapoda: Brachyura: Tanidromitidae) from the Oxfordian of Poland: Acta Zoologica Cracoviensia, 58, 129-136. https://doi.org/10.3409/azc.58_2.129
- Starzyk, N., 2020, The oldest species of the genus Laeviprosopon (Decapoda: Brachyura: Prosopidae) from the Oxfordian of Poland: Acta Zoologica Cracoviensia, 63, 9-13. <https://doi.org/10.3409/azc.63.02>
- Starzyk, N., Krzemińska, E., Krzemiński, W., 2012, A new crab species from the Oxfordian of Poland (Decapoda: Brachyura: Goniodromitidae): Neues Jahrbuch für Geologie und Paläontologie Abhandlungen, 263, 143-153. <https://doi.org/10.1127/0077-7749/2012/0219>
- Von Meyer, H., 1835, Briefliche Mittheilungen, an Prof. Bronn gerichtet: Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde, 325.
- Von Meyer, H., 1842, Ueber die in dem dichten gelben Jurakalk von Aalen in Würtemberg vorkommenden Species des Crustaceengenus Prosopon: Beiträge zur Petrefaktenkunde, 5, 70-75.
- Von Meyer, H., 1857, Briefliche Mittheilungen: Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde, 1857, 556.
- Von Meyer, H., 1860, Die Prosoponiden oder die Familie der Maskenkrebse: Palaeontographica, 7, 183-222.
- Wehner, G., 1988, Über die Prosoponiden (Crustacea, Decapoda): Doctoral thesis, Ludwig-Maximilians-Universität zu München, Germany.
- Zeiss, A., 1964, Zur Malm Gamma-Delta-Grenze in Franken: Geologische Blätter für NO-Bayern und angrenzende Gebiete, 14, 104-115.